



Newly discovered methane consumers in lakes

June 7, 2017 | Andri Bryner

Topics: Biodiversity | Climate Change & Energy | Ecosystems

In freshwater lakes, large amounts of the greenhouse gas methane are oxidized by bacteria before it can be emitted into the atmosphere. A study in Lakes Rotsee and Zug has now shown that the bulk of this work is done, not by “classical” methane consumers, but by filamentous bacteria previously known only as contaminants of water supplies.


When decaying organic matter sinks to the bottom of a lake or ocean, methane is produced as the biomass is broken down. Some of the methane is released from the surface into the atmosphere, where it acts as a potent greenhouse gas, while some is broken down by microorganisms in the water column. In a study just published in the ISME (International Society for Microbial Ecology) Journal, an international research group reports that, as well as the “classical” methane consumers, filamentous bacteria of the genus *Crenothrix* – previously little studied in the environment – are also involved in the methane removal process.

Chance discovery

The role played by *Crenothrix* bacteria was discovered by chance, when the researchers were seeking to quantify methane removal in the Rotsee (Canton Lucerne) and Lake Zug with the aid of stable isotope labelling. In this method, methane molecules are labelled with heavy carbon-13 atoms; when ¹³C-labelled methane is assimilated by bacteria, the individual cells can be visualized by means of imaging mass spectrometry. “Typically,” says co-author Jana Milucka of the Max Planck Institute in Bremen, “these are small round or rod-shaped cells.” In this case, however, the bacteria enriched with ¹³C were not only single round cells but also long, filamentous varieties (see photo): This finding was very surprising, because at that point, the researchers didn’t know these filamentous bacteria could occur so abundantly in nature. “We then started wondering about their role in the environmental

removal of methane”, says Milucka.


Commonly known as well-thread



Eawag geologist Carsten Schubert – an expert on microbial degradation of methane in water – was also surprised by the group’s findings in the two Central Swiss lakes. While the large filamentous *Crenothrix* bacteria have long been known, they are only familiar as a contaminant in drinking water systems (hence their name “well-thread”), where their proliferation can cause clogging of pipes, sand filters and screens. *Crenothrix* bacteria have not previously been detected in lake water, because they have not been specifically sought and they are difficult to identify with molecular genetic methods. Schubert concludes: “We seem to have completely underestimated their role in the biogeochemical cycle.” The researchers have now shown that *Crenothrix* are not only a stable part of the microbial community in freshwater lakes but may also be the most important methane consumers.

Original publication

Kirsten Oswald, Jon S Graf et al.: *Crenothrix* are major methane consumers in stratified lakes; ISME Journal (2017) 00, 1–17. <http://dx.doi.org/10.1038/ismej.2017.77>



*Micrographs showing the methane oxidizers discovered in Lake Rotsee. Left: The “classical” bacteria responsible for methane uptake are small and round. Right: ^{13}C labelling combined with mass spectrometry reveals that methane is also assimilated by much larger, filamentous *Crenothrix*. The small blue (left) and white-outlined structures (right) are non-methanotrophic bacteria.*

Related Files

[This media release as pdf \[pdf, 51 KB\]](#)



Kirsten Oswald
+41 44 633 85 75
kirsten.oswald@eawag.ch

Contact



Carsten Schubert
Tel. +41 58 765 2195
carsten.schubert@eawag.ch



Andri Bryner

Media officer

Tel. +41 58 765 5104

andri.bryner@eawag.ch

<https://www.eawag.ch/en/info/portal/news/news-archive/archive-detail/newly-discovered-methane-consumers-in-lakes>