



## The underground as a haven for biodiversity

October 12, 2021 | Claudia Carle

Topics: Biodiversity


**Europe has relatively low biodiversity compared to most other continents because many species became extinct during the ice ages. In subterranean ecosystems, however, which were shielded from climatic turbulences, a great diversity of ancient species were able to survive. This is the conclusion of a study on the amphipod genus *Niphargus*.**

Around 30 million years ago, numerous mountain ranges began to form in south-eastern Europe as a result of the collision of the European, Adriatic and African tectonic plates. Where these consisted of limestone – such as in the south-eastern Alps, the Carpathians and the Dinaric Alps – caves developed in the rock at the same time under the influence of precipitation. When living organisms colonise newly created habitats such as these caves and find sufficient ecological resources there, a rapid emergence of new species can occur: the immigrated parent species fans out into numerous new species that specialise in the various niches within the new habitat. This basic mechanism of evolution is called “adaptive radiation” and is responsible for much of the biodiversity on Earth. In Europe, however, during the Pleistocene period with its constant alternation of cold and warm periods, a large part of the biodiversity that developed in this way became extinct or migrated to other regions. Therefore, in terms of biodiversity, Europe occupies the lowest ranks among the continents in most species groups. Only Antarctica has fewer species.

### Habitats sealed off from climate chaos

Now, a study which was recently published in the journal “Nature Communications” by Ole Seehausen, Head of the Fish Ecology & Evolution Department at Eawag, and a team of scientists from the University of Ljubljana and the Agricultural Institute of Slovenia, provides evidence that a large proportion of ancient biodiversity has been able to survive in subterranean habitats sealed off from climate chaos. Previously, there were only isolated indications for this. Seehausen and his research


colleagues have now been able to substantiate this assumption for the amphipod genus *Niphargus*. Amphipods are crustaceans of a few millimetres to a few centimetres in size that live in bodies of water. The genus *Niphargus* is only found underground, in cave water or in groundwater. “*Niphargus* is the most species-rich of the amphipod genus and includes hundreds of species whose origins we have studied using various evolutionary genetic modelling techniques,” explains Seehausen. “The results show that this biodiversity developed 15 million years ago, exactly when and exactly where new subterranean habitats formed with the caves in the limestone mountain ranges.” This correlation suggests that colonisation of the new subterranean habitats is the trigger for the fanning out of this amphipod genus into an enormous number of new species.



Numerous new species of *Niphargus* amphipods evolved from one parent species, each with a different size and shape and inhabiting different subterranean habitats and niches. (Photos: Denis Copila?-Ciocianu, Teo Deli?)

### Pollution of the groundwater as the greatest danger

Thus, ironically, such an inhospitable habitat as underground, where there are few nutrients and no light, seems to be a haven of biodiversity in Europe. If their hypothesis can be generalised, the researchers believe that unknown ancient species should also be found among other cave-dwelling organisms. Whether these will ever be discovered, however, also depends on how long they can defy the encroachment of their habitats by humans. Today, the greatest danger to subterranean biodiversity, which has survived all the ice ages, comes from groundwater pollution.



The larger the circle, the greater the number of species in the corresponding group. The groups are named and numbered according to the mountain regions in which they occur:

1. Pontic
2. Pannonian
3. North Dinaric
4. South Dinaric
5. West Balkan
- and 6. the Apennine group.

Cover picture: Denis Copila?-Ciocianu, Teo Deli?

### Original publication

Borko, Š.; Trontelj, P.; Seehausen, O.; Moškri?, A.; Fišer, C. (2021) A subterranean adaptive radiation of amphipods in Europe, *Nature Communications*, 12, 3688 (12 pp.), [doi:10.1038/s41467-021-24023-w](https://doi.org/10.1038/s41467-021-24023-w), [Institutional Repository](#)

### Funding / Partnerships

Slovenian Research Agency University of Ljubljana (Cene Fišer, Špela Borko) Agricultural Institute of Slovenia

### Contact



**Ole Seehausen**

Tel. +41 58 765 2121

[ole.seehausen@eawag.ch](mailto:ole.seehausen@eawag.ch)



**Claudia Carle**

Science editor

Tel. +41 58 765 5946

[claudia.carle@eawag.ch](mailto:claudia.carle@eawag.ch)

<https://www.eawag.ch/en/info/portal/news/news-archive/archive-detail/the-underground-as-a-haven-for-biodiversity>