



## Selenium deficiency promoted by climate change

February 21, 2017 | Martina Schürmann

Topics: Society | Water & Development

**As a result of climate change, concentrations of the trace element selenium in soils are likely to decrease. Because the selenium content of crops may also be reduced, the risk of selenium deficiency could be increased in many regions of the world. This was shown by a recent study which used data-mining to model the global distribution of selenium.**

Selenium is an essential micronutrient obtained from dietary sources such as cereals. The selenium content of foodstuffs largely depends on concentrations in the soil: previous studies have shown that low selenium concentrations are associated with high pH and oxygen availability and low clay and soil organic carbon content. In Europe, as is known from regional studies, selenium-poor soils are found particularly in Germany, Denmark, Scotland, Finland and certain Balkan countries.

To date, however, little has been known about the global distribution of selenium. Using data-mining techniques – involving the assessment of datasets originally collected for other purposes – scientists from Eawag and five other research institutes\* have now modelled global soil selenium concentrations. The sixteen datasets assessed (1994–2016) comprised a total of 33,241 soil data points. Analysis of selenium concentrations in the top 30 centimetres of soil, together with 26 environmental variables, indicated the dominant role of climate-soil interactions in controlling soil selenium distributions.

The main factors influencing soil selenium concentrations are precipitation and the so-called aridity index (potential evaporation:precipitation ratio). Precipitation leads to leaching of selenium from the soil. At the same time, precipitation can have a positive effect on selenium concentrations since the oxygen content is lower in wet than in dry soils, which means that the selenium is less soluble and thus less mobile. In addition, frequent precipitation leads to a low soil pH, which promotes the binding of negatively charged selenium to soil particles. Higher selenium concentrations are most likely to occur in areas with low to moderate precipitation and high clay content, while lower concentrations are found in


arid areas with high pH and low clay content.

In the light of these findings, the scientists modelled mean soil selenium concentrations for the periods 1980–1999 and 2080–2099. Under a moderate climate change scenario, selenium levels are predicted to increase in parts of Australia, China, India and Africa. Overall, however, selenium levels are expected to decrease: by the end of this century (2080–2099), 66% of croplands are predicted to lose selenium (mean decrease of almost 9% compared to 1980–1999). Particularly affected are agricultural areas of Europe and India, China, southern South America, southern Africa and the south-western United States (see map).

These losses could have implications for human health – at present, up to 1 billion people are thought to be affected by low dietary selenium intake. According to the authors, this study serves as an early warning for humanitarian organizations and the agro-industry. Fertilizers containing selenium could be used to combat selenium deficiency, as has been done in Finland since 1984. In addition, selenium additives could be used in animal feed.

### Photos

Download free use only in connection with this media information, no archiving.



*Changes in soil selenium concentrations (percentage increase or decrease) predicted to occur as a result of climate change.*

### Importance of selenium for health

Selenium (Se) is essential for human health and has to be obtained from dietary sources. As an antioxidant, it scavenges free radicals, thus supporting the immune system. It is also required for the synthesis of numerous proteins. Up to 1 billion people are thought to be affected by low dietary selenium intake, which can cause cardiomyopathy. However, excessive intake of selenium can also be harmful, leading to vomiting, diarrhoea and liver damage.



The trace element selenium is taken up from the soil by crops and thus enters the food chain.  
Photo: Agroscope (Gabriela Brändle, Urs Zihlmann), LANAT (Andreas Chervet)

\* Eawag: Swiss Federal Institute of Aquatic Science and Technology  
ETH Zurich (Institute of Atmospheric and Climate Science and Institute of Biogeochemistry  
and Pollutant Dynamics)  
Potsdam Institute for Climate Impact Research  
Rothamsted Research  
University of Aberdeen

#### **Publication original**

Jones, G. D.; Droz, B.; Greve, P.; Gottschalk, P.; Poffet, D.; McGrath, S. P.; Seneviratne, S. I.;  
Smith, P.; Winkel, L. H. E. (2017) Selenium deficiency risk predicted to increase under future  
climate change, *Proceedings of the National Academy of Sciences of the United States of  
America PNAS*, 114(11), 2848-2853, [doi:10.1073/pnas.1611576114](https://doi.org/10.1073/pnas.1611576114), [Institutional Repository](#)

## **Related Files**

[This media release as pdf](#) [pdf, 271 KB]

## **Project video (3:13)**





## Contact



**Lenny Winkel**

Tel. +41 58 765 5601

[lenny.winkel@eawag.ch](mailto:lenny.winkel@eawag.ch)



**Andri Bryner**

Media officer

Tel. +41 58 765 5104

[andri.bryner@eawag.ch](mailto:andri.bryner@eawag.ch)

<https://www.eawag.ch/en/info/portal/news/news-archive/archive-detail/selenium-deficiency-promoted-by-climate-change>