



## Climate change leads to disproportionately high pollutant absorption

May 10, 2023 | Cornelia Zogg

Topics: Biodiversity | Ecosystems | Pollutants | Climate Change & Energy

**Pesticides and other chemicals harm biodiversity, and climate change is also threatening many habitats. With his study, Eawag researcher Johannes Rath has shown that these two challenges for our society may even worsen. Amphipods absorbed pollutants faster as water temperatures rose. The researcher has received an award from the Society of Environmental Toxicology and Chemistry for his paper on the toxicokinetics of amphipods in combination with increased water temperatures.**

Environmental pollution and climate change are two of the greatest threats to ecosystems and biodiversity. We know a great deal about these two factors, but little research has been done into how they interact. Several studies in recent years have concluded that native invertebrates in aquatic environments are more sensitive to pollutants when water temperatures rise. In his study, Eawag researcher Johannes Rath looked into the question of whether this increased sensitivity can be explained by changes in toxicokinetic processes.

### Higher temperatures lead to increased pollutant absorption

Together with researchers from the ETH, the Carinthia University of Applied Sciences and the Helmholtz Centre for Environmental Research, Rath studied the biochemical aspects of amphipods at four different temperatures and with twelve polar organic pollutants. These small aquatic animals are a fundamental component of the aquatic food web and are practically native all over the world.

Rath's research revealed a clear connection between toxicokinetic rates and water temperature. Toxicokinetic rates describe how pollutants are processed in the body, i.e. the resorption, distribution,

metabolism and excretion of the pollutants. This means: as water temperatures increase, amphipods absorb, metabolise and excrete more pollutants, not just linearly but also exponentially. Rath was also able to show that the influence of temperature on toxicokinetic rates was on a similar scale to the influence of temperature on physiological rates, for example, the animals' respiration. This connection can simplify the practical application of the results in models of pollutant absorption.



**The warmer the water, the faster amphipods absorb pollutants.**  
(Photo: Johannes Rath, Eawag)

The findings show that dependencies exist between environmental pollution and climate change and that these can become stronger. According to Rath: "High temperatures and peaks of pollutant concentration in the water often occur together, for example following pesticide application in spring and summer. This makes the temperature-related acceleration of toxicokinetic processes especially worrying, as much higher peak concentrations can be achieved in the organisms than previously assumed." However, more studies, including with other substance classes, are necessary to obtain an even better picture of the underlying mechanisms. The results give new insights into how aquatic life absorbs chemicals in different climate scenarios. These results can therefore contribute to improving assessment of environmental risks.

### **Award for publication**

Johannes Rath has just received the SETAC Europe Rifcon Early Career Scientist Award for the paper. The award for early-stage researchers was set up to support members of the Society of Environmental Toxicology and Chemistry (SETAC) in their career development. It is conferred for original scientific research, political or other professional achievements by an early-stage researcher. The Eawag researcher was presented with this prestigious award at the ceremony last Sunday.

Cover picture: Eawag researcher Johannes Rath (right) has won the SETAC Europe Rifcon Early Career Scientist Award for his paper on the toxicokinetics of amphipods at increased water temperatures. Here, receiving the award in Dublin. (Photo: SETAC)

### **Original publication**



decoration:none;font-style:italic}.extbase-debugger-center .extbase-debug-property{color:#f1f1f1}.extbase-debugger-center .extbase-debug-closure{color:#9BA223;}Extbase Variable Dumparray(3 items) publications => '26212' (5 chars) libraryUrl => '' (0 chars) layout => '0' (1 chars) Extbase Variable Dumparray(1 item) 0 => Snowflake\Publications\Domain\Model\Publicationprototypepersistent entity (uid=26212, pid=124) originalId => protected26212 (integer) authors => protected'Raths,&nbsp;J.; Švara,&nbsp;V.; Lauper,&nbsp;B.; Fu,&nbsp;Q.; Hollender,&nbsp;sp;J.' (81 chars) title => protected'Speed it up: how temperature drives toxicokinetics of organic contaminants i n freshwater amphipods' (98 chars) journal => protected'Global Change Biology' (21 chars) year => protected2023 (integer) volume => protected29 (integer) issue => protected'5' (1 chars) startpage => protected'1390' (4 chars) otherpage => protected'1406' (4 chars) categories => protected'aquatic invertebrates; Arrhenius; bioconcentration; biotransformation; Gamma rus pulex; Hyalella azteca; micropollutants' (119 chars) description => protected'The acceleration of global climate change draws increasing attention towards interactive effects of temperature and organic contaminants. Many studies reported a higher sensitivity of aquatic invertebrates towards contaminant exposure with increasing or fluctuating temperatures. The hypothesis of this study was that the higher sensitivity of invertebrates is associated with the changes of toxicokinetic processes that determine internal concentrations of contaminants and consequently toxic effects. Therefore, the influence of temperature on toxicokinetic processes and the underlying mechanisms were studied in two key amphipod species (<em>Gammarus pulex</em> and <em>Hyalella azteca</em>). Bioconcentration experiments were carried out at four different temperatures with a mixture of 12 exposure relevant polar organic contaminants. Tissue and medium samples were taken in regular intervals and analysed by online solid-phase extraction liquid chromatography high-resolution tandem mass spectrometry. Subsequently, toxicokinetic rates were modelled and analysed in dependence of the exposure temperature using the Arrhenius equation. An exponential relationship between toxicokinetic rates versus temperature was observed and could be well depicted by applying the Arrhenius equation. Due to a similar Arrhenius temperature of uptake and elimination rates, the bioconcentration factors of the contaminants were generally constant across the temperature range. Furthermore, the Arrhenius temperature of the toxicokinetic rates and respiration was mostly similar. However, in some cases (citalopram, cyprodinil), the bioconcentration factor appeared to be temperature dependent, which could potentially be explained by the influence of temperature on active uptake mechanisms or biotransformation. The observed temperature effects on toxicokinetics may be particularly relevant in non-equilibrated systems, such as exposure peaks in summer as exemplified by the exposure modelling of a field me...)' (2302 chars) serialnumber => protected'1354-1013' (9 chars) doi => protected'10.1111/gcb.16542' (17 chars) uid => protected26212 (integer) \_localizedUid => protected26212 (integer)modified \_languageUid => protectedNULL \_versionedUid => protected26212 (integer)modified pid => protected124 (integer) Raths, J.; Švara, V.; Lauper, B.; Fu, Q.; Hollender, J. (2023) Speed it up: how temperature drives toxicokinetics of organic contaminants in freshwater amphipods, *Global Change Biology*, 29(5), 1390-1406, doi:10.1111/gcb.16542, [Institutional Repository](#)

## Related Links

Website SETAC

## Contact



**Johannes Raths**

Postdoctoral scientist (he/him)

Tel. +41 58 765 5739

[johannes.raths@eawag.ch](mailto:johannes.raths@eawag.ch)



**Juliane Hollender**

Senior scientist / Group leader

Tel. +41 58 765 5493

[juliane.hollender@eawag.ch](mailto:juliane.hollender@eawag.ch)



**Cornelia Zogg**

Science Editor

Tel. +41 58 765 5763

[cornelia.zogg@eawag.ch](mailto:cornelia.zogg@eawag.ch)

<https://www.eawag.ch/en/info/portal/news/news-archive/archive-detail/climate-change-leads-to-disproportionately-high-pollutant-absorption>