



## Underground microbial life more stable than expected

July 1, 2025 | Universität Texas / Eawag

Topics: Ecosystems

**Microbial communities living deep underground in the groundwater of Lavey-les-Bains remain surprisingly stable throughout the year, even though the composition of the water changes seasonally. This is the finding of a study by Eawag researchers that contributes to a better understanding of geothermal systems and microbial life deep underground.**

Thermal groundwater systems, such as those found in the Alps near Lavey-les-Bains, are naturally warm, underground bodies of water, some of which emerge as hot springs on the earth's surface. These systems are rich in gases and minerals. However, they also harbour ecosystems in which microbes exist without sunlight by using sulphur, iron or hydrogen for their metabolism. Nevertheless, it is difficult to study these microbial communities as their habitats are virtually inaccessible. For this reason, there is still only limited knowledge of thermal groundwater systems, which are also interesting because of their potential for climate-friendly energy production.

A study recently published in the journal "Proceedings of the National Academy of Sciences (PNAS)" now sheds light on these hidden ecosystems. In connection with his doctoral thesis, Sébastien Giroud from the Water Resources & Drinking Water department at the aquatic research institute Eawag investigated how microbes in the subsurface respond to seasonal changes in the chemical composition of the water. Giroud's team thus sampled the groundwater at depths of between 200 and 500 metres all year round. Dissolved ions, water isotopes and noble gases were measured in order to record variability in the water chemistry. DNA sequencing also provided information on changes in microbial composition.



One of the measuring points in Lavey-les-Bains, where thermal water is pumped from a depth of 200 metres. The mobile gas analyser (“miniRuedi”) in the orange case analyses the gases contained in the thermal water.

(Photo: Sébastien Giroud, Eawag)

The findings are unexpected: although geochemical indicators such as electrical conductivity and the concentration of dissolved gases show clear seasonal fluctuations in the composition of the water, the microbial communities remain stable. The varying composition of the water is due to the seasonal differences in the mixture of younger, near-surface and old groundwater from the depths within the thermal system. This is mainly caused by snowmelt and the associated groundwater recharge in summer. The microbial communities, whose composition varies with depth, hardly change over the course of the year. For example, sulphur-degrading bacteria are prevalent at depths of 200 metres, while bacteria that utilise sulphate, iron or hydrogen predominate at depths of 500 metres.

These results suggest that it is not the water composition but other environmental conditions, in this case temperature, that determine microbial life in deep continental groundwater systems. The data also shows the remarkable resilience of microbial life deep in the earth. “This qualifies the assumption that all life underground is influenced by the conditions on the surface,” says Sébastien Giroud. However, according to the researcher, it is all the more important to investigate the consequences of interventions underground – such as the extraction or supply of thermal energy (see also box).



```

TFIMFYwaDExVjExeiBNMTAsMUgxdjloOVYxeilvPjxnIGkPSJJbm5lcil+PHJIY3QgeD0iMilgeT
0iNSIgy2xhc3M9InNOMClgd2lkdGg9ljlGhlaWdodD0iMSlvpjwvZz48L3N2Zz4=)}.extbase-
debugger{display:block;text-align:left;background:#2a2a2a;border:1px solid #2a2a2a;box-
shadow:0 3px 0 rgba(0,0,0,.5);color:#000;margin:20px;overflow:hidden;border-radius:4px}.ext
base-debugger-floating{position:relative;z-index:999}.extbase-debugger-
top{background:#444;font-size:12px;font-family:monospace;color:#f1f1f1;padding:6px
15px}.extbase-debugger-center{padding:0 15px;margin:15px 0;background-image:repeating-
linear-gradient(to bottom,transparent 0,transparent 20px,#252525 20px,#252525
40px)}.extbase-debugger-center,.extbase-debugger-center .extbase-debug-string,.extbase-
debugger-center a,.extbase-debugger-center p,.extbase-debugger-center pre,.extbase-
debugger-center strong{font-size:12px;font-weight:400;font-family:monospace;line-
height:20px;color:#f1f1f1}.extbase-debugger-center pre{background-color:transparent;margin:
0;padding:0;border:0;word-wrap:break-word;color:#999}.extbase-debugger-center .extbase-
debug-string{color:#ce9178;white-space:normal}.extbase-debugger-center .extbase-debug-
type{color:#569CD6;padding-right:4px}.extbase-debugger-center .extbase-debug-
unregistered{background-color:#dce1e8}.extbase-debugger-center .extbase-debug-
filtered,.extbase-debugger-center .extbase-debug-proxy,.extbase-debugger-center .extbase-
debug-ptype,.extbase-debugger-center .extbase-debug-visibility,.extbase-debugger-center
.extbase-debug-scope{color:#fff;font-size:10px;line-height:12px;padding:2px 4px;margin-
right:2px;position:relative;top:-1px}.extbase-debugger-center .extbase-debug-
scope{background-color:#497AA2}.extbase-debugger-center .extbase-debug-
ptype{background-color:#698747}.extbase-debugger-center .extbase-debug-
visibility{background-color:#698747}.extbase-debugger-center .extbase-debug-
dirty{background-color:#FFFFFFB6}.extbase-debugger-center .extbase-debug-
filtered{background-color:#4F4F4F}.extbase-debugger-center .extbase-debug-seeabove{text-
decoration:none;font-style:italic}.extbase-debugger-center .extbase-debug-
property{color:#f1f1f1}.extbase-debugger-center .extbase-debug-
closure{color:#9BA223;}Extbase Variable Dumparray(3 items) publications => '34784' (5
chars) libraryUrl => " (0 chars) layout => '0' (1 chars) Extbase Variable Dumparray(1 item) 0
=> Snowflake\Publications\Domain\Model\Publicationprototypepersistent entity (uid=34784,
pid=124) originalId => protected34784 (integer) authors => protected'Giroud,&nbsp;S.;
Deng,&nbsp;L.; Lever,&nbsp;M.&nbsp;A.; Schilling,&nbsp;O.&n
bsp;S.; Kipfer,&nbsp;R.' (99 chars) title => protected'Resilience of deep aquifer microbial
communities to seasonal hydrological fl
uctuations' (86 chars) journal => protected'Proceedings of the National Academy of
Sciences of the United States of Amer
ica PNAS' (84 chars) year => protected2025 (integer) volume => protected122 (integer)
issue => protected'23' (2 chars) startpage => protected'e2422608122 (9 pp.)' (19 chars)
otherpage => protected" (0 chars) categories => protected'thermal aquifer; deep microbial
communities; noble gases; geomicrobiological' (76 chars) description => protected'The
influence of seasonal variations in temperature and precipitation on sub
surface biogeochemical processes remains poorly understood. In the Lavey-les
-Bains thermal system in the Swiss Alps, annual variations in electrical con
ductivity are observed to depths of 500 m, suggesting a potential link to su
rface environmental changes. Here we show, through year-round analyses of st
able water isotopes, noble gases, and conductivity, that seasonally varying
contributions of shallow groundwater from the Rhône alluvial aquifer mix wi
th deep groundwater. Despite vertically similar fluid geochemical compositio

```


ns suggesting high hydrological connectivity, microbial communities exhibit significant depth-dependent variation with minimal seasonal change. This decoupling of dynamic water source partitioning and stable microbial community structure has not been previously observed and fills a critical gap in our understanding of geothermal systems and microbial life in the deep subsurface. At 200 m, the communities are dominated by sulfur-disproportionating Bacteria (*Dissulfurispira*) and Micrarchaeota, while at 500 m the major groups include sulfate- and iron-reducers and/or hydrogen-oxidizers (Thermals, Thermodesulfobacteriota, and Bathyarchaeota). Our study highlights the resilience of terrestrial subsurface microbial communities to temporal variations in water sources and fluid composition. We propose that intrinsic environmental properties - such as temperature - are more critical drivers of microbial community structure in hydrologically connected deep aquifers than seasonal hydrological changes.' (1623 chars) serialnumber => protected'0027-8424' (9 chars) doi => protected'10.1073/pnas.2422608122' (23 chars) uid => protected34784 (integer) \_localizedUid => protected34784 (integer)modified \_languageUid => protectedNULL \_versionedUid => protected34784 (integer)modified pid => protected124 (integer) Giroud, S.; Deng, L.; Lever, M. A.; Schilling, O. S.; Kipfer, R. (2025) Resilience of deep aquifer microbial communities to seasonal hydrological fluctuations, *Proceedings of the National Academy of Sciences of the United States of America PNAS*, 122(23), e2422608122 (9 pp.), doi:10.1073/pnas.2422608122, [Institutional Repository](#)

### Funding / Partnerships

Eawag ETH Zurich University of Basel University of Texas at Austin Canton of Valais

[Project website](#)

### Contact



**Sébastien Giroud**

Tel. +41 58 765 5388

[sebastien.giroud@eawag.ch](mailto:sebastien.giroud@eawag.ch)



**Oliver Schilling**

Group Leader Tracer Hydrogeology

Tel. +41 58 765 5931

[oliver.schilling@eawag.ch](mailto:oliver.schilling@eawag.ch)



**Rolf Kipfer**

Tel. +41 58 765 5530  
[rolf.kipfer@eawag.ch](mailto:rolf.kipfer@eawag.ch)



**Claudia Carle**

Science editor  
Tel. +41 58 765 5946  
[claudia.carle@eawag.ch](mailto:claudia.carle@eawag.ch)

<https://www.eawag.ch/en/info/portal/news/news-archive/archive-detail/underground-microbial-life-more-stable-than-expected>