



Alpine fish biodiversity is amazingly young

October 7, 2022 | Bärbel Zierl

Topics: Biodiversity

A high fraction of the endemic biodiversity of the Alps is very old. The endemics – species found only in a confined area – have developed over the past millions of years during the cycles of glacial and interglacial periods or even before these cycles began. Fish, however, are an exception: most endemic fish species emerged only after the end of the last ice age about 15,000 years ago.

The Alps are a habitat for numerous plants and animals – a hotspot of biodiversity on the European continent. As part of the Blue-Green Biodiversity research initiative of the Eawag and WSL research institutes, a group of researchers has now investigated how and when this diversity arose, both on land and in the water. They analysed 497 alpine and perialpine species – from around the Alps – belonging to five main groups: 121 fish, 39 amphipods, 31 amphibians, 158 butterflies and 148 flowering plants.

The analyses show that the Ice Ages have significantly shaped today's alpine biodiversity. Since the Pleistocene, which began about 2.6 million years ago and ended about 12,000 years ago, several warm and cold periods alternated. During the cold periods, glaciers advanced into the valleys and changed the habitats of flora and fauna. Only small ice-free areas far away from each other on the mountain peaks or in the lowlands offered opportunities for retreat for the displaced plants and animals. Due to the spatial separation of populations, they evolved differently, giving rise to many new species.



Evolutionary and ecological history of a hypothetical biodiversity in an alpine-like system. (Graphic: Liz Amman, modified from Jardim de Queiroz et al.)

Large differences in endemic species

Particularly interesting for the researchers were the endemic species – species that are only found in a very narrow geographic area. Luiz Jardim de Queiroz, researcher at the aquatic research institute Eawag and the University of Bern as well as lead author of the study recently published in the scientific journal *Proceedings of the Royal Society B: Biological Sciences*, says: "While the endemic species of amphipods, amphibians, butterflies and flowering plants are largely very old and originated during or even before the Pleistocene, most of the endemic fish species are surprisingly young. The vast majority of them only appeared after the end of the last glacial period about 15,000 years ago."


Since many, if not all, alpine and perialpine waters were covered by a thick layer of ice during the Pleistocene Ice Ages, freshwater habitats were unsuitable for fish species, and local fish populations most probably died out in the Alps. Only after the end of the last Ice Age were they able to migrate again likely from lower-lying river sections of the large rivers such as the Danube, the Rhine or the Rhone. "In the perialpine lakes, a few of these colonisers then specialised within a short time, adapting to several ecological niches in the same lakes and thus formed several new species in most lakes, all of which endemic," says Luiz Jardim de Queiroz.

The amphipods examined in this study did not suffer the same fate as the fish. Since unlike fish they do not depend on large, open water habitats, they were able to survive the Ice Ages in small outlets or under the glaciers in caves and in groundwater. Therefore, most of the endemic amphipod species of the Alps are ancient and some even predate the Pleistocene. The amphibians, in turn, need free water for breeding only in spring and summer and spend the rest of their lives on land. They therefore also found refugia in the Alps during the Ice Ages. And as for the terrestrial species, many of the butterflies and plant species found refugial habitats to persist through the Ice Ages too.

The fish are slow, but creative colonisers

The researchers suggest that the difference in the pace of speciation between fish and the other groups is due to how fast they could colonise the new habitats that opened up after the last Ice Age. Terrestrial species should have colonised those areas relatively fast, as they can

disperse in any direction. Butterflies, for example, fly from place to place without facing great obstacles. The seeds of many plants, in turn, can be transported by the wind or by birds over great distances.



Two endemic lake sculpins from Lake Thun: the littoral Aare-lakes sculpin (dark with large protruding eyes) is endemic to lakes in the Aare system, the profundal Lake Thun sculpin that is endemic to Lake Thun. (Photo: Ole Seehausen)

Fish species, however, must first navigate through widely branching rivers to colonise the ecologically highly isolated lakes that emerged in the headwaters of the river networks after the glaciers retreated. This takes time, especially for the species that came from more distant temperate or warm water refugia. Most river fish also found the deep and cold lakes had little to offer them. "Therefore, only a few newcomers, adapted to cold water, such as whitefish and charrs, that probably arrived early from relatively nearby Pleistocene lowland areas, were able to occupy this habitat and diversify," adds Ole Seehausen, department head at Eawag, professor at the University of Bern and senior author of the study.

This different speed of dispersal influences the pace of speciation. If new habitats are quickly colonised by many different newcomers, all available niches are occupied within a short time. That is what probably occurred in the terrestrial alpine habitat. Therefore, plants and animals have fewer opportunities to evolve new specialisations and form new species. The first cold water fish species to migrate back to the alpine lakes, however, found diverse, unused habitats and were able to diversify rapidly. The researchers see this as one of the main reasons for the astonishingly fast pace of speciation in some fish after the end of the last Ice Age.

The researchers hope that the new findings will also contribute to the protection of alpine biodiversity. "If we understand how biodiversity evolved in the past, we can better foresee how species will respond to future climate changes, allowing us to develop strategies to preserve them," says Luiz Jardim de Queiroz.

This study is part of the research project "[Biodiversity assembly in blue and green ecosystems: speciation versus immigration](#)", which contributes to the [Blue-Green Biodiversity research initiative](#) – an Eawag-WSL collaboration that addresses biodiversity across aquatic


```
.extbase-debug-scope{color:#fff;font-size:10px;line-height:12px;padding:2px 4px;margin-right:2px;position:relative;top:-1px}.extbase-debugger-center .extbase-debug-scope{background-color:#497AA2}.extbase-debugger-center .extbase-debug-type{background-color:#698747}.extbase-debugger-center .extbase-debug-visibility{background-color:#698747}.extbase-debugger-center .extbase-debug-dirty{background-color:#FFFFB6}.extbase-debugger-center .extbase-debug-filtered{background-color:#4F4F4F}.extbase-debugger-center .extbase-debug-seeabove{text-decoration:none;font-style:italic}.extbase-debugger-center .extbase-debug-property{color:#f1f1f1}.extbase-debugger-center .extbase-debug-closure{color:#9BA223;}Extbase Variable Dumparray(2 items) publications => '25593' (5 chars) libraryUrl => '' (0 chars) Extbase Variable Dumparray(1 item) 0 => Snowflake\Publications\Domain\Model\Publicationprototypepersistent entity (uid=25593, pid=124) originalId => protected25593 (integer) authors => protected'Jardim de Queiroz,&nbsp;L.; Doenz,&nbsp;C.&nbsp;J.; Altermatt,&nbsp;F.; Alther,&nbsp;R.; Borko,&nbsp;Š.; Brodersen,&nbsp;J.; Gossner,&nbsp;M.&nbsp;M.; Graham,&nbsp;C.; Matthews,&nbsp;B.; McFadden,&nbsp;I.&nbsp;R.; Pellissier,&nbsp;L.; Schmitt,&nbsp;T.; Selz,&nbsp;O.&nbsp;M.; Villalba,&nbsp;S.; Rüber,&nbsp;L.; Zimmermann,&nbsp;N.&nbsp;E.; Seehausen,&nbsp;O.' (360 chars) title => protected'Climate, immigration and speciation shape terrestrial and aquatic biodiversity in the European Alps' (99 chars) journal => protected'Proceedings of the Royal Society B: Biological Sciences' (55 chars) year => protected2022 (integer) volume => protected289 (integer) issue => protected'1980' (4 chars) startpage => protected'20221020 (10 pp.)' (17 chars) otherpage => protected'' (0 chars) categories => protected'time for speciation; allopatric speciation; adaptive radiation; Pleistocene refugia; glacial species pump; European Alps' (120 chars) description => protected'Quaternary climate fluctuations can affect speciation in regional biodiversity assembly in two non-mutually exclusive ways: a glacial species pump, where isolation in glacial refugia accelerates allopatric speciation, and adaptive radiation in underused adaptive zones during ice-free periods. We detected biogeographic and genetic signatures associated with both mechanisms in the assembly of the biota of the European Alps. Age distributions of endemic and widespread species within aquatic and terrestrial taxa (amphipods, fishes, amphibians, butterflies and flowering plants) revealed that endemic fish evolved only in lakes, are highly sympatric, and mainly of Holocene age, consistent with adaptive radiation. Endemic amphipods are ancient, suggesting preglacial radiation with limited range expansion and local Pleistocene survival, perhaps facilitated by a groundwater-dwelling lifestyle. Terrestrial endemics are mostly of Pleistocene age and are thus more consistent with the glacial species pump. The lack of evidence for Holocene adaptive radiation in the terrestrial biome is consistent with faster recolonization through range expansion of these taxa after glacial retreats. More stable and less seasonal ecological conditions in lakes during the Holocene may also have contributed to Holocene speciation in lakes. The high proportion of young, endemic species makes the Alpine biota vulnerable to climate change, but the mechanisms and consequences of species loss will likely differ between biomes because of their distinct evolutionary histories.' (1563 chars) serialnumber => protected'0962-8452' (9 chars) doi => protected'10.1098/rspb.2022.1020' (22 chars) uid => protected25593 (integer) _localizedUid => protected25593 (integer)modified _languageUid =>
```

protectedNULL _versionedUid => protected25593 (integer)modified pid => protected124 (integer) Jardim de Queiroz, L.; Doenz, C. J.; Altermatt, F.; Alther, R.; Borko, Š.; Brodersen, J.; Gossner, M. M.; Graham, C.; Matthews, B.; McFadden, I. R.; Pellissier, L.; Schmitt, T.; Selz, O. M.; Villalba, S.; Rüber, L.; Zimmermann, N. E.; Seehausen, O. (2022) Climate, immigration and speciation shape terrestrial and aquatic biodiversity in the European Alps, *Proceedings of the Royal Society B: Biological Sciences*, 289(1980), 20221020 (10 pp.), doi:10.1098/rspb.2022.1020, [Institutional Repository](#)

Contact WSL

[Prof. Dr. Niklaus Zimmermann](#)
Tel. +41 44 739 23 37

Contact



Ole Seehausen

Tel. +41 58 765 2121

ole.seehausen@eawag.ch



Bärbel Zierl

Science editor

Tel. +41 58 765 6840

baerbel.zierl@eawag.ch

<https://www.eawag.ch/en/info/portal/news/news-archive/archive-detail/alpine-fish-biodiversity-is-amazingly-young>